

Spectral cutoffs in indirect dark matter searches

Christoph Weniger*

Max-Planck-Institut für Physik, Föhringer Ring 6, 80805 Munich, Germany

E-mail: weniger@mppmu.mpg.de

Torsten Bringmann¹, Francesca Calore¹ and Gilles Vertongen²

¹ II. Institute for Theoretical Physics, University of Hamburg, Luruper Chaussee 149, DE-22761 Hamburg, Germany

² Institut d'Astrophysique de Paris, UMR-7095 du CNRS, 98 bis bd Arago, 75014 Paris, France

* presenting author

Abstract. Indirect searches for dark matter annihilation or decay products in the cosmic-ray spectrum are plagued by the question of how to disentangle a dark matter signal from the omnipresent astrophysical background. One of the practically background-free smoking-gun signatures for dark matter would be the observation of a sharp cutoff or a pronounced bump in the gamma-ray energy spectrum. Such features are generically produced in many dark matter models by internal Bremsstrahlung, and they can be treated in a similar manner as the traditionally looked-for gamma-ray lines. Here, we discuss prospects for seeing such features with present and future Atmospheric Cherenkov Telescopes.

1. Introduction

Indirect dark matter (DM) searches aim at seeing an excess in cosmic rays from the annihilation or decay of DM in the Galactic halo [1]. Very often, indirect searches focus on *secondary photons* from the fragmentation of hadronic annihilation products. The corresponding spectra are rather broad and peak at energies much lower than the DM mass m_χ , which generically makes a convincing claim of a DM detection above the astrophysical backgrounds difficult. In many models, however, *pronounced spectral features* are expected at the kinematic endpoint $E_\gamma = m_\chi$; they include monochromatic gamma-ray lines [2], sharp steps or cutoffs [3, 4] as well as pronounced bumps [5]. The type and strength of these features are intricately linked to the particle nature of DM; a detection would thus not only allow a convincing discrimination from astrophysical backgrounds but also to determine important DM model parameters like the value of m_χ . So far, only line-signals have explicitly been searched for [6]—despite the fact that they are loop-suppressed and thus generically subdominant compared to other spectral signatures [5]. Here, we discuss a general method to search for sharp spectral features in gamma-ray observations. Concentrating on DM models with a large internal Bremsstrahlung (IB) component, and on observations with Imaging Atmospheric Cherenkov Telescopes (IACTs), we derive projected limits and prospects to see such signatures with current and future instruments.

	DM particle	m_χ^{th} [TeV]	$\langle\sigma v\rangle^{\text{th}}$ [cm ³ s ⁻¹]	relevant channel	spectral feature
$\gamma\gamma$	any WIMP	$\mathcal{O}(0.1\text{--}10)$	$\mathcal{O}(10^{-30})$	$\gamma\gamma$	line
KK	$B^{(1)}$	1.3	$1 \cdot 10^{-26}$	$\ell^+\ell^-\gamma$	FSR step
BM3	neutralino	0.23	$9 \cdot 10^{-29}$	$\ell^+\ell^-\gamma$	IB bump
BM4	neutralino	1.9	$3 \cdot 10^{-27}$	$W^+W^-\gamma$	IB bump

Table 1. DM benchmark models used in our analysis as examples for the typical spectral endpoint features to be expected in WIMP annihilations. For these particular models, we also state the annihilation channel that is most important in this context, as well as mass and total annihilation rate for thermally produced DM. See text and Ref. [10] for further details about the DM models and Fig. 1 for the corresponding photon spectra.




Figure 1. Photon spectra for the DM benchmark models of Tab. 1. Dashed lines show the same spectra smeared with a Gaussian of width $\Delta x/x = 0.1$. From Ref. [10].

2. Methods and Targets

The defining aspect of the above-mentioned spectral features is an abrupt change of the gamma-ray flux as function of energy. It is therefore possible to concentrate the search for spectral features on a small sliding energy window $[E_0, E_1]$, with $E_0 < m_\chi < E_1$, and window sizes of the order of a few times the energy resolution of the instrument. An important advantage of considering only small window sizes is that astrophysical gamma-ray fluxes can be typically very well described by a simple power-law. Limits on and the significance for dark matter induced fluxes can then be derived by standard statistical methods like the profile likelihood method.

We are interested in deriving projected limits on the dark matter annihilation cross-section that follow from the spectral end-point features of different dark matter scenarios *alone*, *i.e.* without taking into account secondary photons, using the above method. We will focus on observations of the Galactic center region with IACTs, considering benchmark scenarios that roughly correspond to the telescope characteristics of the currently operating H.E.S.S. (IACT1) [7], the future CTA (IACT2) [8] and—as the most optimistic choice for indirect DM searches—the proposed Dark Matter Array (DMA, IACT3) [9]. For the background, we take into account cosmic-ray fluxes of electrons and protons, the diffuse gamma-ray flux and the source HESS J1745-290 at (or very close to) the Galactic center. We adopt an Einasto dark matter profile, and a relatively small target region $\Delta\Omega = 2^\circ \times 2^\circ$ around the Galactic center. The energy window size is chosen such that the impact of a nonzero background curvature on our final results is less than 50% (see Ref. [10] for details).

We will discuss three types of typical endpoint features that arise from radiative corrections to the tree-level annihilation process: (1) The most striking spectral signature, in terms of a




Figure 2. *Thick lines:* Expected 2σ upper limits on $\langle\sigma v\rangle$ for selected DM models, DM profiles and observational scenarios; bands indicate the variance of these limits. *Thin lines:* Spectral feature of DM signal has $\mathcal{S}/\mathcal{B} \approx 1\%$ (after convolution with energy dispersion). The *left panel* shows limits on gamma-ray lines, rescaled by a loop-factor of α_{em}^{-2} for better comparison. In the *central panel*, the gray band indicates the expected $\langle\sigma v\rangle$ for KK DM, the black part being compatible with the observed relic density. In the *right panel*, we indicate the adopted neutralino benchmark points, and the *dotted lines* show the projected 5σ sensitivity. The benchmark experiments IACT1, IACT2 and IACT3 correspond roughly to H.E.S.S., CTA and DMA, respectively, see text for details. The figure is taken from Ref. [10].

possible discrimination from a power-law background, is a *gamma-ray line*, which would result from the direct annihilation of DM into $\gamma\gamma$, $Z\gamma$ or $H\gamma$. Generically, the annihilation rate is loop-suppressed and expected to be of the order of $\langle\sigma v\rangle_{\text{line}} \sim \alpha_{\text{em}}^2 \times \langle\sigma v\rangle_{\text{tree}} \sim 10^{-30} \text{cm}^3 \text{s}^{-1}$ (although in some cases much stronger line signals are possible). (2) As an example for a step-like feature we use the gamma-ray spectrum expected from annihilating *Kaluza-Klein (KK) DM* in models of universal extra dimensions [11]. Its total gamma-ray annihilation spectrum at high energies is dominated by final state radiation (FSR) off lepton final states, and the shape of the spectrum dN/dx , with $x = E_\gamma/m_\chi$, turns out to be essentially independent of m_χ and other model parameters [4]. (3) Pronounced bump-like features at $E \simeq m_\chi$ may arise from IB in the annihilation of *neutralino DM* [5]. Here, BM3 is a typical example for a neutralino in the stau co-annihilation region, where photon emission from virtual sleptons greatly enhances the photon spectrum at high energies; BM4 refers to a situation in which IB from $W^\pm$ final states dominates.

In Tab. 1, we shortly summarize the properties of the DM benchmark models described above, including for completeness the actual DM mass and total annihilation rate needed to obtain the observed relic density for thermally produced DM. Note, however, that we essentially treat these values as free parameters in our analysis and that we are rather interested in the spectral shape of the annihilation signal, represented by dN/dx ; in Fig. 1 we show these spectra for a direct comparison.

3. Results and Discussion

In Fig. 2 we show our results for the expected 2σ upper limits (thick lines) on the above DM models as well as the variance of these limits among the mock data sets. We find that in particular IB features in the spectrum (right panel) have the potential to constrain the annihilation rate at least down to values typically expected for thermal production, $\langle\sigma v\rangle \sim$

$3 \cdot 10^{-26} \text{cm}^3 \text{s}^{-1}$, already for modest assumptions about the DM distribution. For models with very large IB contributions like BM3, we find that our method would provide even stronger limits on $\langle\sigma v\rangle$ than what was obtained by the HESS analysis of the Galactic center region assuming annihilation into $\bar{b}b$ [12].

For the case of not too strongly pronounced endpoint features (like line signals in most models or the step for Kaluza-Klein DM), secondary photons will usually be more powerful in constraining the total annihilation rate $\langle\sigma v\rangle$. However, in case of an adiabatically compressed profile our limits could improve by maybe two orders of magnitude, as demonstrated for gamma-ray lines in the left panel. As shown in the central panel of Fig. 2, the future CTA should be able to place limits about one order of magnitude stronger than currently possible, and the proposed DMA could further improve these by another factor of ten.

Finally, we would like to stress that our limits in general provide rather *complementary information* on the DM nature and can thus usually not easily be compared with limits on secondary photons. In any case, from the point of view of indirect DM searches, the *detection* of the kinematic cutoff will be much more interesting than the detection of secondary photons, because they would provide rather unambiguous evidence for the DM nature of the signal as well as allow to determine important parameters like the DM mass. An obvious extension of the approach presented here is to apply it to the *discrimination* of models [13].

Acknowledgments

T.B. and C.W. thank the organizers of the TAUP 2011 for an interesting and stimulating conference. T.B. and F.C. acknowledge support from the German Research Foundation (DFG) through Emmy Noether grant BR 3954/1-1.

- [1] G. Bertone, D. Hooper, J. Silk, Phys. Rept. **405** (2005) 279.
- [2] L. Bergström, P. Ullio and J. H. Buckley, Astropart. Phys. **9** (1998) 137 [arXiv:astro-ph/9712318].
- [3] A. Birkedal, K. T. Matchev, M. Perelstein and A. Spray, arXiv:hep-ph/0507194.
- [4] L. Bergström, T. Bringmann, M. Eriksson and M. Gustafsson, Phys. Rev. Lett. **94** (2005) 131301 [arXiv:astro-ph/0410359].
- [5] T. Bringmann, L. Bergström and J. Edsjö, JHEP **0801** (2008) 049 [arXiv:0710.3169 [hep-ph]].
- [6] A.A. Pullen, R.R. Chary, M. Kamionkowski, Phys. Rev. D **76** (2007) 063006 [Erratum-ibid. **83** (2011) 029904] [arXiv:astro-ph/0610295]; F. A. Aharonian *et al.* [HEGRA Collaboration], Astron. Astrophys. **400**, 153 (2003) [astro-ph/0302347]; A. A. Abdo *et al.*, Phys. Rev. Lett. **104** (2010) 091302 [arXiv:1001.4836 [astro-ph.HE]]; G. Vertongen and C. Weniger, JCAP **1105** (2011) 027 [arXiv:1101.2610 [hep-ph]].
- [7] F. Aharonian *et al.* [H.E.S.S. Collaboration], Astron. Astrophys. **457** (2006) 899 [arXiv:astro-ph/0607333].
- [8] The CTA Consortium, arXiv:1008.3703 [astro-ph.IM].
- [9] L. Bergström, T. Bringmann and J. Edsjö, Phys. Rev. D **83** (2011) 045024 [arXiv:1011.4514 [hep-ph]].
- [10] T. Bringmann, F. Calore, G. Vertongen and C. Weniger, Phys. Rev. D **84** (2011) 103525 [arXiv:1106.1874 [hep-ph]].
- [11] G. Servant and T. M. P. Tait, Nucl. Phys. B **650** (2003) 391 [arXiv:hep-ph/0206071].
- [12] A. Abramowski *et al.*, Phys. Rev. Lett. **106**, 161301 (2011) [arXiv:1103.3266 [astro-ph.HE]].
- [13] T. Bringmann, F. Calore, G. Vertongen and C. Weniger, work in progress; see also, *e.g.*, M. Perelstein and B. Shakya, JCAP **1010** (2010) 016 [arXiv:1007.0018 [astro-ph.HE]].